

Star Scheduling

Nadia Brauner* Hadrien Cambazard* Benoit Cance*
Nicolas Catusse* Pierre Lemaire* Bernard Penz*
Anne-Marie Lagrange^o Pascal Rubini^o

* Grenoble Alpes, CNRS, G-SCOP, F-38000 Grenoble
pierre.lemaire@grenoble-inp.fr

^o CNRS, IPAG, F-38000 Grenoble, France

june 2015

Star Scheduling


Nadia Brauner* Hadrien Cambazard* Benoit Cance*
Nicolas Catusse* Pierre Lemaire* Bernard Penz*
Anne-Marie Lagrange^o Pascal Rubini^o

* Grenoble Alpes, CNRS, G-SCOP, F-38000 Grenoble
pierre.lemaire@grenoble-inp.fr

^o CNRS, IPAG, F-38000 Grenoble, France

june 2015

Looking at the stars




- $[r_s; d_s)$ is the visibility interval
- p_s is the required duration of observation
- w_s is the interest

scheduling the observation of star s means observing s for a continuous duration p_s within the visibility interval $[r_s; d_s)$, rewarding w_s


Star scheduling (one night)

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s)$



Star scheduling (one night)

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s]$




Question: find $\mathcal{S}' \subset \mathcal{S}$ and starting times $t_s, \forall s \in \mathcal{S}'$ such that

- for each $s \in \mathcal{S}'$: $[t_s; t_s + p_s] \subset [r_s; d_s]$
- for each $(s_1, s_2) \in \mathcal{S}'^2$: $[t_{s_1}; t_{s_1} + p_{s_1}] \cap [t_{s_2}; t_{s_2} + p_{s_2}] = \emptyset$
- $\sum_{s \in \mathcal{S}'} w_s$ is maximized


Star scheduling (several nights)

Instance: a set \mathcal{N} of nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$, depending on the night n of the observation




Star scheduling (several nights)

Instance: a set \mathcal{N} of nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$, depending on the night n of the observation



The order is known!

The meridian instant ($m_s = \frac{r_s + d_s}{2}$) is a **mandatory** instant of observation, that is: for every star s , $p_s^n \geq \frac{d_s^n - r_s^n}{2}$




Property

If for all star s : $p_s^n \geq \frac{d_s^n - r_s^n}{2}$, then observations must be scheduled by non-decreasing meridian time

The order is known!

The meridian instant ($m_s = \frac{r_s + d_s}{2}$) is a **mandatory** instant of observation, that is: for every star s , $p_s^n \geq \frac{d_s^n - r_s^n}{2}$




Property

If for all star s : $p_s^n \geq \frac{d_s^n - r_s^n}{2}$, then observations must be scheduled by non-decreasing meridian time

The order is known!

The meridian instant ($m_s = \frac{r_s + d_s}{2}$) is a **mandatory** instant of observation, that is: for every star s , $p_s^n \geq \frac{d_s^n - r_s^n}{2}$



Property

If for all star s : $p_s^n \geq \frac{d_s^n - r_s^n}{2}$, then observations must be scheduled by non-decreasing meridian time

A MIP model

$$\max \sum_{s \in \mathcal{S}} w_s z_s \longrightarrow = 1 \text{ iff } s \text{ observed}$$

$$\text{s.c. } \sum_{n \in \mathcal{N}} z_s^n = z_s \longrightarrow = 1 \text{ iff } s \text{ observed on night } n$$

A MIP model

$$\max \sum_{s \in \mathcal{S}} w_s z_s \longrightarrow = 1 \text{ iff } s \text{ observed}$$

$$s.c. \sum_{n \in \mathcal{N}} z_s^n = z_s \longrightarrow = 1 \text{ iff } s \text{ observed on night } n$$

$$\text{visibility window of night } n \left\{ \begin{array}{l} r_s^n z_s^n \leq t_s \longrightarrow \text{starting time of observation } s \\ t_s + p_s^n z_s^n \leq d_s^n z_s^n + M(1 - z_s^n) \end{array} \right.$$

A MIP model

$$\max \sum_{s \in \mathcal{S}} w_s z_s \longrightarrow = 1 \text{ iff } s \text{ observed}$$

$$s.c. \sum_{n \in \mathcal{N}} z_s^n = z_s \longrightarrow = 1 \text{ iff } s \text{ observed on night } n$$

$$\text{visibility window of night } n \left\{ \begin{array}{l} r_s^n z_s^n \leq t_s \longrightarrow \text{starting time of observation } s \\ t_s + p_s^n z_s^n \leq d_s^n z_s^n + M(1 - z_s^n) \end{array} \right.$$

$$s \prec s' \text{ if observed the same night} \left\{ \begin{array}{l} z_s^n + z_{s'}^n - 1 \leq y_{ss'} \longrightarrow = 1 \text{ iff } s \text{ and } s' \text{ observed the same night} \\ t_s + p_s^n \leq t_{s'} + M(1 - z_{ss'}) \end{array} \right.$$

Introduction

Complexity

Solving star scheduling

Conclusions and perspectives

Scheduling one night

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s)$ such that $p_s \geq (d_s - r_s)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

Scheduling one night

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s)$ such that $p_s \geq (d_s - r_s)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $1|r_j| \sum w_j U_j$ is NP-complete (Lenstra et al., 1977)

Scheduling one night

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s)$ such that $p_s \geq (d_s - r_s)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $1|r_j| \sum w_j U_j$ is NP-complete (Lenstra et al., 1977)
- $1|r_j, p_j = p| \sum w_j U_j$ is polynomial (Baptiste, 1999)

Scheduling one night

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s)$ such that $p_s \geq (d_s - r_s)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $1|r_j| \sum w_j U_j$ is NP-complete (Lenstra et al., 1977)
- $1|r_j, p_j = p| \sum w_j U_j$ is polynomial (Baptiste, 1999)
- what if $p_j \geq (d_j - r_j)/2$?

Scheduling one night

Instance: a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s and a visibility window $[r_s; d_s)$ such that $p_s \geq (d_s - r_s)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $1|r_j| \sum w_j U_j$ is NP-complete (Lenstra et al., 1977)
- $1|r_j, p_j = p| \sum w_j U_j$ is polynomial (Baptiste, 1999)
- what if $p_j \geq (d_j - r_j)/2$?

Complexity of the one night case

Star scheduling of one night is NP-Hard (even if $w_s = 1$)

Scheduling of one night is NP-hard

- Variant of Partition : $2n$ pairs (a_{2i-1}, a_{2i}) so that $\sum_i a_i = 2B$.
Can a total of B be made with one item from each pair?

Scheduling of one night is NP-hard


- Variant of Partition : $2n$ pairs (a_{2i-1}, a_{2i}) so that $\sum_i a_i = 2B$.
Can a total of B be made with one item from each pair?
- For each pair of items (a_{2i}, a_{2i+1}) create a pair of incompatible stars (s_{2i-1}, s_{2i}) : same visibility window and $p_s \geq (d_s - r_s)/2$



$$p_{2i-1} = p_{2i} = 2B + a_i$$


Scheduling of one night is NP-hard

- Variant of Partition : $2n$ pairs (a_{2i-1}, a_{2i}) so that $\sum_i a_i = 2B$.
Can a total of B be made with one item from each pair?
- For each pair of items (a_{2i}, a_{2i+1}) create a pair of incompatible stars (s_{2i-1}, s_{2i}) : same visibility window and $p_s \geq (d_s - r_s)/2$
- Set visibility windows so that scheduling a star from a pair does not prevent scheduling a star from another pair and each pair of stars have dedicated instants



Scheduling of one night is NP-hard

- Variant of Partition : $2n$ pairs (a_{2i-1}, a_{2i}) so that $\sum_i a_i = 2B$.
Can a total of B be made with one item from each pair?
- For each pair of items (a_{2i}, a_{2i+1}) create a pair of incompatible stars (s_{2i-1}, s_{2i}) : same visibility window and $p_s \geq (d_s - r_s)/2$
- Set visibility windows so that scheduling a star from a pair does not prevent scheduling a star from another pair and each pair of stars have dedicated instants
- “yes” to Partition \iff non-idling schedule of length $(2n + 1)B$



A pseudo-polynomial algorithm

$f(i, t)$: maximum interest with stars 1 to i , and such that s_i ends before time t

$f(i, t) =$

$$\begin{cases} \min(f(i-1, t), f(i-1, t - p_i) + w_i) & \forall i \in [1, m], t \in [r_i + p_i, T] \\ f(i-1, t) & \forall i \in [1, m], t \in [0, r_i + p_i[\\ -\infty & \forall i \in [1, m], t < 0 \\ 0 & i = 0, \forall t \in [0, T] \end{cases}$$

We are looking for $f(m, T)$ which can be computed in $O(mT)$

Scheduling several nights

Instance: a set of n nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$ such that $p_s^n \geq (d_s^n - r_s^n)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

Scheduling several nights

Instance: a set of n nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$ such that $p_s^n \geq (d_s^n - r_s^n)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $P|r_j| \sum w_j U_j$ (identical nights) is NP-complete

Scheduling several nights

Instance: a set of n nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$ such that $p_s^n \geq (d_s^n - r_s^n)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $P|r_j| \sum w_j U_j$ (identical nights) is NP-complete
- $p_s = (d_s - r_s)$, identical nights: polynomial (Kolen et al., 2007)

Scheduling several nights

Instance: a set of n nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$ such that $p_s^n \geq (d_s^n - r_s^n)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $P|r_j| \sum w_j U_j$ (identical nights) is NP-complete
- $p_s = (d_s - r_s)$, identical nights: polynomial (Kolen et al., 2007)
- what if $p_j \geq (d_j - r_j)/2$?

Scheduling several nights

Instance: a set of n nights, a set \mathcal{S} of stars; each star $s \in \mathcal{S}$ has an interest w_s , an observation duration p_s^n and a visibility window $[r_s^n; d_s^n)$ such that $p_s^n \geq (d_s^n - r_s^n)/2$; a bound W

Question: find a subset of stars so that the total interest is at least W , visibility windows are respected and observations do not overlap

- $P|r_j| \sum w_j U_j$ (identical nights) is NP-complete
- $p_s = (d_s - r_s)$, identical nights: polynomial (Kolen et al., 2007)
- what if $p_j \geq (d_j - r_j)/2$?

Complexity of the several nights case

Star scheduling of several nights is unary NP-Hard (even if $w_s = 1$ and all nights are identical)

Logic-based Bender decomposition

Master problem:
assignment of stars to nights

$$\begin{array}{ll}\max & \sum_{s \in \mathcal{S}} w_s z_s \\ \text{s.c.} & \sum_{n \in \mathcal{N}} z_s^n = z_s\end{array}$$

Logic-based Bender decomposition

Master problem:
assignment of stars to nights

Slave problem:
scheduling of each nights

$$\begin{array}{ll} \max & \sum_{s \in \mathcal{S}} w_s z_s \\ \text{s.c.} & \sum_{n \in \mathcal{N}} z_s^n = z_s \end{array}$$


Night 1: ok?


Night 2: ok?

...

Logic-based Bender decomposition

Master problem:
assignment of stars to nights

Slave problem:
scheduling of each nights

$$\begin{aligned}
 \max \quad & \sum_{s \in \mathcal{S}} w_s z_s \\
 \text{s.c.} \quad & \sum_{n \in \mathcal{N}} z_s^n = z_s \\
 & \sum_{s \in C_k^n} z_s^n \leq |C_k^n| - 1
 \end{aligned}$$


Night 1: ok?


Night 2: ok?

...

Logic-based Bender decomposition

Master problem:
assignment of stars to nights

Slave problem:
scheduling of each nights

$$\begin{aligned}
 \max \quad & \sum_{s \in \mathcal{S}} w_s z_s \\
 \text{s.c.} \quad & \sum_{n \in \mathcal{N}} z_s^n = z_s \\
 & \sum_{s \in C_k^n} z_s^n \leq |C_k^n| - 1
 \end{aligned}$$


Night 1: ok?

Night 2: ok?

...

- efficient MIP
- upper bound at each iteration
- n independent problems
- linear complexity

Column generation

Night patterns: Ω_n , set of all possible schedules for night n

Pattern k for night n : $p_{n1}^k \dots p_{n|S|}^k$, where $p_{ns}^k = 1$ iff star s belongs to the k -th pattern of night n ; weight $w_n^k = \sum_{s \in S} w_s p_{ns}^k$

Column generation

Night patterns: Ω_n , set of all possible schedules for night n

Pattern k for night n : $p_{n1}^k \dots p_{n|S|}^k$, where $p_{ns}^k = 1$ iff star s belongs to the k -th pattern of night n ; weight $w_n^k = \sum_{s \in S} w_s p_{ns}^k$

$$\begin{aligned} \max \sum_{n \in \mathcal{N}} \sum_{k \in \Omega_n} w_n^k \rho_n^k & \longrightarrow = 1 \text{ iff pattern } k \text{ used, night } n \\ \sum_{k \in \Omega_n} \rho_n^k & = 1 \quad \forall n \in \mathcal{N} \\ \sum_{n \in \mathcal{N}} \sum_{k \in \Omega_n} p_{ns}^k \rho_n^k & \leq 1 \quad \forall s \in S \\ \rho_n^k & \in \{0, 1\} \quad \forall n \in \mathcal{N}, \forall k \in \Omega_n \end{aligned}$$

Column generation

Night patterns: Ω_n , set of all possible schedules for night n

Pattern k for night n : $p_{n1}^k \dots p_{n|S|}^k$, where $p_{ns}^k = 1$ iff star s belongs to the k -th pattern of night n ; weight $w_n^k = \sum_{s \in S} w_s p_{ns}^k$

$$\begin{array}{ll}
 \max_{\alpha_n} & \sum_{n \in \mathcal{N}} \sum_{k \in \Omega_n} w_n^k \rho_n^k \longrightarrow = 1 \text{ iff pattern } k \text{ used, night } n \\
 \beta_s & \sum_{n \in \mathcal{N}} \sum_{k \in \Omega_n} p_{ns}^k \rho_n^k \leq 1 \quad \forall s \in S \\
 & \rho_n^k \in \{0, 1\} \quad \forall n \in \mathcal{N}, \forall k \in \Omega_n
 \end{array}$$

Reduced cost of ρ_n^k : $w_n^k - \alpha_n - \sum_{s \in S} p_{ns}^k \beta_s$

Column generation

Night patterns: Ω_n , set of all possible schedules for night n

Pattern k for night n : $p_{n1}^k \dots p_{n|S|}^k$, where $p_{ns}^k = 1$ iff star s belongs to the k -th pattern of night n ; weight $w_n^k = \sum_{s \in S} w_s p_{ns}^k$

$$\begin{array}{ll}
 \max_{\alpha_n} & \sum_{n \in \mathcal{N}} \sum_{k \in \Omega_n} w_n^k \rho_n^k \longrightarrow = 1 \text{ iff pattern } k \text{ used, night } n \\
 \beta_s & \sum_{n \in \mathcal{N}} \sum_{k \in \Omega_n} p_{ns}^k \rho_n^k \leq 1 \quad \forall s \in S \\
 & \rho_n^k \in \{0, 1\} \quad \forall n \in \mathcal{N}, \forall k \in \Omega_n
 \end{array}$$

Reduced cost of ρ_n^k : $w_n^k - \alpha_n - \sum_{s \in S} p_{ns}^k \beta_s$

Pattern of max reduced cost? Single Night Case with costs $w_s - \beta_s$

Local search

Classical local-search procedure

Neighborhoods:

- moving a star from one night to another
- inserting an unobserved star
- exchanging two stars

An optimal schedule for each night is computed systematically

Solutions

| instance | $ S $ | $ N $ | BD (OPT/ <i>UB</i>) | | CG (UB) | | LS (LB) | |
|----------|-------|-------|----------------------|--------|--------------|--------|--------------|--------|
| | | | val | cpu(s) | val | cpu(s) | val | cpu(s) |
| pb1 | 200 | 32 | 5200 | 900 | 5200 | 1.47 | 5200 | 0.28 |
| pb2 | 200 | 32 | 3310 | 900 | 3310 | 0.99 | 3310 | 0.33 |
| pb3 | 200 | 69 | 7800 | 100 | 7800 | 1.59 | 7800 | 0.17 |
| pb4 | 200 | 69 | - | - | 4870 | 1.63 | 4870 | 0.11 |
| pb5 | 400 | 69 | 12660 | 900 | 11910 | 5.11 | 11910 | 12.39 |
| pb6 | 400 | 69 | 9250 | 900 | 9099.9 | 19.57 | 9070 | 773.95 |
| pb7 | 400 | 142 | - | - | 13680 | 11.85 | 13680 | 0.15 |
| pb8 | 400 | 142 | 9760 | 900 | 9760 | 13.71 | 9760 | 0.21 |
| real | 800 | 142 | 18930 | 900 | 18620 | 92.41 | 18510 | 689.60 |
| | | | | | | | 18480 | 306.07 |

Example solution

solution

Star scheduling

- a particular interval-scheduling problem with known order
- NP-hardness is proven for both one night and general cases
- several solution methods are proposed and tested

Work in progress...

- Instances
 - we need more instances!
 - study the structure of the real instance (e.g., similarity between nights)
- Solution methods
 - enrich numerical experiments
 - embed the CG into a branch & bound to get optimal solutions
 - analyze (and improve) LS behavior
 - get advantage of nights' similarities
- Complexity
 - draw precise complexity frontiers
 - study approximability